arXiv:1303.6669v1 [math.MG] 26 Mar 2013

ISHIKAWA ITERATION PROCESS IN CAT(K) SPACES

C. JUN

ABSTRACT. In this paper, we establish Δ-convergence results for Ishikawa iterations in complete CAT(K) spaces.

1. INTRODUCTION

Let (X, d) be a complete metric space and let T be a mapping from X to X. Then T is called *nonexpansive* if for all $x, y \in X$,

$$d(T(x), T(y)) \leq d(x, y).$$

A point $x \in X$ is called a *fixed point* of T if $T(x) = x$. $\mathrm{Fix}(T)$ denotes the set of fixed points of T. Kirk proved the existence of fixed points for nonexpansive mappings on CAT(0) spaces in [6] and CAT(K) spaces in [7].

A CAT(K) space is a metric space in which no triangle is fatter than the triangle with the same edge lengths in a model space, which is the 2-dimensional, complete, simply-connected space of constant curvature K (see Definition 1). A CAT(K) space is a generalization of a simply-connected Riemannian manifold with sectional curvature $\leq K$; we will introduce generalized definitions of convergence and notations including the sum $\oplus$, which interpolates between a pair of points along a geodesic, in Section 2.

In a CAT(K) space X, for $t_n, s_n \in [0, 1]$ and $x_0 \in X$, the Ishikawa iteration $\{x_n\}$ is defined by

$$x_{n+1} = t_n T(y_n) \oplus (1 - t_n)x_n, \quad n \geq 0, \tag{1.1}$$

where $y_n = s_n T(x_n) \oplus (1 - s_n)x_n$.

For a nonexpansive mapping T, Dhompongsa and Panyanak in [2] obtained a Δ-convergence result for Ishikawa iterations in complete CAT(0) spaces under the conditions

$$\sum_{n=0}^{\infty} t_n(1 - t_n) = \infty, \sum_{n=0}^{\infty} (1 - t_n)s_n < \infty \ \textit{ and } \ \limsup_n s_n < 1.$$

In [9], a similar result was proved by Panyanak and Laokul under the other conditions

$$\sum_{n=0}^{\infty} t_n(1 - t_n) = \infty \ \textit{ and } \ \sum_{n=0}^{\infty} t_n(1 - t_n)s_n < \infty.$$

In this paper, we will obtain Δ-convergence results for Ishikawa iterations in complete CAT(K) spaces (see Theorem 16 and Theorem 19).

1991 *Mathematics Subject Classification.* 47H09, 53C20.

Key words and phrases. Ishikawa iteration, CAT(K) geometry, Nonexpansive mapping.

In [5], He, Fang, López and Li studied the Δ-convergence of Mann iterations in complete $\mathrm{CAT}(K)$ spaces with the condition

$$\sum_{n=0}^{\infty} t_n(1-t_n)=\infty.$$

Since the Mann iteration is given by (1.1) when $s_n=0$ for all n, we provide an alternative proof of the Δ-convergence theorem for Mann iterations in complete $\mathrm{CAT}(K)$ spaces.

2. PRELIMINARIES

Let (X,d) be a metric space. The open ball centered at p with radius r is denoted by $B_r(p)$. The closed ball centered at p with radius r is denoted by $B_r[p]$.

A curve $\gamma : I \to X$ is called a *geodesic* if for any two $t,t' \in I$, $d(\gamma(t),\gamma(t')) = |t-t'|$. We denote by $[xy]$, a unit-speed geodesic $\gamma : I \to X$ from x to y defined on $I=[0,t]$, where $\gamma(0)=x$, $\gamma(t)=y$ and $t=d(x,y)$. By $\triangle xyz$, we denote the geodesic triangle of geodesics $[xy]$,$[xz]$ and $[yz]$.

Let C be a positive constant. A metric space X is a *geodesic space* if any two points are joined by a geodesic; and a *C-geodesic space* if any two points with distance $< C$ are joined by a geodesic. A set $Y \subset X$ is *convex* if any two points $x,y \in Y$ can be joined by a geodesic and all geodesics joining them are contained in Y. If this condition holds for any two points in Y with distance $< C$, Y is said to be *C-convex*.

For a constant K, we use M_K to denote the 2-dimensional, complete, simply-connected space of constant curvature K. Then $M_0=\mathbb{E}^2$, $M_1=\mathbb{S}^2$ and $M_{-1}=\mathbb{H}^2$. Let d_K be the metric of M_K. D_K denotes the diameter of M_K. Thus, $D_K=\frac{\pi}{\sqrt{K}}$ if $K>0$ and $D_K=\infty$ if $K \le 0$.

A triangle $\triangle \tilde{x}_1\tilde{x}_2\tilde{x}_3$ in M_K is called a *comparison triangle* for $\triangle x_1x_2x_3$ in X if $d_K(\tilde{x}_i,\tilde{x}_j)=d(x_i,x_j)$ for $i,j \in \{1,2,3\}$.

Definition 1. Let (X,d) be a metric space and let K be a real constant. A D_K-geodesic space X is a $\mathrm{CAT}(K)$ *space* if for any geodesic triangle $\triangle xy_1y_2$ of perimeter $< 2D_K$, and its comparison triangle $\triangle \tilde{x}\tilde{y}_1\tilde{y}_2$ in M_K, we have

$$d(z_1,z_2) \le d_K(\tilde{z}_1,\tilde{z}_2),$$

where z_i is any point on $[xy_i]$ and $\tilde{z}_i$ is the point on $[\tilde{x}\tilde{y}_i]$ such that $d_K(\tilde{x},\tilde{z}_i)=d(x,z_i)$ for $i \in \{1,2\}$.

We now record a few lemmas about $\mathrm{CAT}(K)$ spaces that we will need in the sequel.

Lemma 2. [1, Page 160] *Let X be a $\mathrm{CAT}(K)$ space.*
(1) *For any two points x,y in X with distance less than D_K, there is a unique geodesic $[xy]$ connecting them.*
(2) *Any ball in X with radius less than $D_K/2$ is convex.*

Lemma 3. [1, Page 178] *Let (X,d) be a $\mathrm{CAT}(K)$ space and let F be a closed and D_K-convex subset of X. Then for each point $x \in X$ such that $d(x,F) < D_K/2$, there is a unique point $y \in F$ such that $d(x,y)=d(x,F)$.*

Let (X,d) be a $\mathrm{CAT}(K)$ space and let $x,y \in X$ such that $d(x,y) < D_K$. Then $tx \oplus (1-t)y$ denotes the unique point on $[xy]$ for $t \in [0,1]$ such that $d(x,tx \oplus (1-t)y)=(1-t)d(x,y)$ and $d(y,tx \oplus (1-t)y)=td(x,y)$.

Lemma 4. [10, Lemma 3.3] *For a positive number* $C \leq \pi/2$*, let* (X, d) *be a* $\mathrm{CAT}(1)$ *space and let* $p, x, y \in X$ *such that* $d(p, x) \leq C$*,* $d(p, y) \leq C$ *and* $d(x, y) \leq C$*. Then for any* $t \in [0, 1]$*,*

$$d((1-t)p \oplus tx, (1-t)p \oplus ty) \leq \frac{\sin tC}{\sin C} d(x, y).$$

We can get the next lemma by following the proof of Prop. 3.1 in [11] with $\varepsilon = \pi/4$.

Lemma 5. *Let* (X, d) *be a* $\mathrm{CAT}(1)$ *space. Then there is a constant* $k > 0$ *such that*

$$d^2(x, ty \oplus (1-t)z) \leq td^2(x, y) + (1-t)d^2(x, z) - \frac{k}{2}t(1-t)d^2(y, z)$$

for any $t \in [0, 1]$ *and any points* $x, y, z \in X$ *such that* $d(x, y) \leq \pi/4$*,* $d(x, z) \leq \pi/4$ *and* $d(y, z) \leq \pi/2$*.*

For a bounded sequence $\{x_n\}$ in X, define

$$r(x, \{x_n\}) = \limsup_{n \to \infty} d(x, x_n).$$

The *asymptotic radius* of $\{x_n\}$ is defined by

$$r(\{x_n\}) = \inf\{r(x, \{x_n\}) : x \in X\}.$$

The *asymptotic center* of $\{x_n\}$ is defined by

$$A(\{x_n\}) = \{x \in X : r(x, \{x_n\}) = r(\{x_n\})\}.$$

Now we can give a definition of Δ-convergence, and list a few properties.

Definition 6. For a bounded sequence $\{x_n\}$ in X, the sequence $\{x_n\}$ is said to *Δ-converge to* $x \in X$ if x is the unique asymptotic center of $\{u_n\}$ for every subsequence $\{u_n\}$ of $\{x_n\}$. In this case, we will write $\Delta - \lim_n x_n = x$ and call x the *Δ-limit of* $\{x_n\}$.

Then we have a lemma to show a property of a sequence which Δ-converges.

Lemma 7. [5, Prop. 2.3] *Let* (X, d) *be a complete* $\mathrm{CAT}(K)$ *space and let* $p \in X$*. Suppose that a sequence* $\{x_n\}$ *Δ-converges to* x *such that* $r(p, \{x_n\}) < D_K/2$*. Then*

$$d(x, p) \leq \liminf_{n \to \infty} d(x_n, p).$$

Definition 8. Let (X, d) be a complete metric space and let F be a nonempty subset of X. Then a sequence $\{x_n\}$ in X is *Fejér monotone* with respect to F if

$$d(x_{n+1}, q) \leq d(x_n, q)$$

for all $n \geq 0$ and all $q \in F$.

Definition 9. [5] For a sequence $\{x_n\}$ in X, a point $x \in X$ is a *Δ-cluster point* of $\{x_n\}$ if there exists a subsequence of $\{x_n\}$ that Δ-converges to x.

With Definition 9, we will know when a sequence $\{x_n\}$ in X Δ-converges to a point of F if $\{x_n\}$ is Fejér monotone with respect to F.

Lemma 10. [5, Lemma 3.2] *Let* X *be a complete* $\mathrm{CAT}(K)$ *space and let* F *be a nonempty subset of* X*. Suppose that the sequence* $\{x_n\}$ *of* X *is Fejér monotone with respect to* F *and the asymptotic radius* $r(\{x_n\})$ *of* $\{x_n\}$ *is less than* $D_K/2$*. If any Δ-cluster point* x *of* $\{x_n\}$ *belongs to* F*, then the sequence* $\{x_n\}$ *Δ-converges to a point of* F*.*

Lemma 11. [12][13] *Suppose that* $\{a_n\}$ *and* $\{b_n\}$ *satisfy that*

$$a_n \geq 0, \ b_n \geq 0 \ \textit{and} \ \ a_{n+1} \leq (1+b_n)a_n$$

for all $n \geq 0$*. If* $\sum_{n=0}^{\infty} b_n$ *converges, then* $\lim_{n\to\infty} a_n$ *exists. Additionally, if there is a subsequence of* $\{a_n\}$ *which converges to* 0*, then* $\lim_{n\to\infty} a_n = 0$*.*

3. ISHIKAWA ITERATION PROCESS ON CAT(K) SPACES

Lemma 12. [10, Theorem 3.4] *Let* X *be a complete* $\mathrm{CAT}(1)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$*. Then* F *is closed and* π*-convex.*

Lemma 13. *Let* X *be a complete* $\mathrm{CAT}(1)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$*. If* $\{x_n\}$ *is defined by* (1.1) *for* $x_0 \in X$ *such that* $d(x_0, F) \leq \pi/4$*, then there is a unique point* p *in* F *such that* x_n*,* y_n*,* $T(x_n)$ *and* $T(y_n)$ *are at distance* $\leq d(x_0, p)$ *from* p*.*

Proof. Since $d(x_0, F) \leq \pi/4$, by Lemma 3 and 12, there is a unique point p in F such that $d(x_0, p) = d(x_0, F)$. By induction, we want to show that

$$d(p, y_n) \leq d(p, x_n) \leq d(p, x_0).$$

For $n = 0$, since T is nonexpansive, we have $d(p, T(x_0)) \leq d(p, x_0) \leq \pi/4$. Since $B_{\pi/4}[p]$ is convex, we get

$$d(p, y_0) = d(p, s_0T(x_0) \oplus (1-s_0)x_0) \leq d(p, x_0).$$

Suppose that $d(p, y_n) \leq d(p, x_n) \leq d(p, x_0)$. Since $d(p, T(y_n)) \leq d(p, y_n) \leq \pi/4$ and $B_{\pi/4}[p]$ is convex,

$$d(p, x_{n+1}) = d(p, t_nT(y_n) \oplus (1-t_n)x_n) \leq d(p, x_n).$$

Since $d(p, T(x_{n+1})) \leq d(p, x_{n+1}) \leq \pi/4$,

$$d(p, y_{n+1}) = d(p, s_{n+1}T(x_{n+1}) \oplus (1-s_{n+1})x_{n+1}) \leq d(p, x_{n+1}).$$

Therefore

$$d(p, y_{n+1}) \leq d(p, x_{n+1}) \leq d(p, x_n).$$

$\square$

We will prove Lemma 14 and Lemma 15, which are obtained by following the proofs in [9].

If $d(x_0, F) = 0$, then $x_0 \in F$ and hence by definition (1.1), $x_n = x_0$ for all n. So we just consider the case $d(x_0, F) > 0$.

Lemma 14. *Let* X *be a complete* $\mathrm{CAT}(1)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$*. If* $\{x_n\}$ *is defined by* (1.1) *for* $x_0 \in X$ *such that* $d(x_0, F) \leq \pi/4$*, then*

$$d(T(x_{n+1}), x_{n+1}) \leq [1 + 4\frac{C}{\sin C}t_n(1-t_n)s_n]d(T(x_n), x_n)$$

for all $n \geq 0$ *where* $C := 2d(x_0, F)$*.*

Proof. Since $d(T(x_n), t_nT(x_n) \oplus (1-t_n)x_n) = (1-t_n)d(T(x_n), x_n)$ and T is nonexpansive, we get

$$\begin{aligned}
d(T(x_{n+1}), x_{n+1}) &\leq d(T(x_{n+1}), T(t_nT(x_n) \oplus (1-t_n)x_n)) \\
&\quad + d(T(t_nT(x_n) \oplus (1-t_n)x_n), T(x_n)) \\
&\quad + d(T(x_n), t_nT(x_n) \oplus (1-t_n)x_n) \\
&\quad + d(t_nT(x_n) \oplus (1-t_n)x_n, x_{n+1}) \\
&\leq 2d(t_nT(x_n) \oplus (1-t_n)x_n, x_{n+1}) \\
&\quad + d(t_nT(x_n) \oplus (1-t_n)x_n, x_n) \\
&\quad + (1-t_n)d(T(x_n), x_n).
\end{aligned}$$

Since $d(t_nT(x_n) \oplus (1-t_n)x_n, x_n) = t_nd(T(x_n), x_n)$, it becomes

$$d(T(x_{n+1}), x_{n+1}) \leq 2d(t_nT(x_n) \oplus (1-t_n)x_n, x_{n+1}) + d(T(x_n), x_n).$$

Note that $d(x_n, T(x_n)) \leq C$, $d(x_n, T(y_n)) \leq C$ and $d(T(x_n), T(y_n)) \leq C$ from Lemma 13. Since $d(x_n, y_n) = s_nd(T(x_n), x_n)$ and $C \leq \pi/2$, by Lemma 4,

$$\begin{aligned}
d(T(x_{n+1}), x_{n+1}) &\leq 2\frac{\sin t_nC}{\sin C}d(T(x_n), T(y_n)) + d(T(x_n), x_n) \\
&\leq 2\frac{t_nC}{\sin C}d(x_n, y_n) + d(T(x_n), x_n) \\
&= (1 + 2\frac{t_nC}{\sin C}s_n)d(T(x_n), x_n).
\end{aligned}$$

Then, multiplying by $(1-t_n)$, we have

$$(1-t_n)d(T(x_{n+1}), x_{n+1}) \leq [1 - t_n + 2\frac{C}{\sin C}t_n(1-t_n)s_n]d(T(x_n), x_n). \tag{3.1}$$

Also, since $d(T(y_n), t_nT(y_n) \oplus (1-t_n)y_n) = (1-t_n)d(T(y_n), y_n)$ and T is nonexpansive, we get

$$\begin{aligned}
d(T(x_{n+1}), x_{n+1}) &\leq d(T(x_{n+1}), T(t_nT(y_n) \oplus (1-t_n)y_n)) \\
&\quad + d(T(t_nT(y_n) \oplus (1-t_n)y_n), T(y_n)) \\
&\quad + d(T(y_n), t_nT(y_n) \oplus (1-t_n)y_n) \\
&\quad + d(t_nT(y_n) \oplus (1-t_n)y_n, x_{n+1}) \\
&\leq 2d(t_nT(y_n) \oplus (1-t_n)y_n, x_{n+1}) \\
&\quad + d(t_nT(y_n) \oplus (1-t_n)y_n, y_n) \\
&\quad + (1-t_n)d(T(y_n), y_n).
\end{aligned}$$

Since $d(t_nT(y_n) \oplus (1-t_n)y_n, y_n) = t_nd(T(y_n), y_n)$, it becomes

$$d(T(x_{n+1}), x_{n+1}) \leq 2d(t_nT(y_n) \oplus (1-t_n)y_n, x_{n+1}) + d(T(y_n), y_n).$$

Since $d(x_n, T(y_n)) \leq C$, $d(y_n, T(y_n)) \leq C$ and $d(x_n, y_n) \leq C$, by Lemma 4,

$$\begin{aligned} d(T(x_{n+1}), x_{n+1}) &\leq 2\frac{\sin(1-t_n)C}{\sin C} d(x_n, y_n) + d(T(y_n), y_n) \\ &\leq 2\frac{(1-t_n)C}{\sin C} d(x_n, y_n) + d(T(y_n), y_n) \\ &\leq 2\frac{(1-t_n)C}{\sin C} d(x_n, y_n) + d(T(y_n), T(x_n)) + d(T(x_n), y_n) \\ &\leq 2\frac{(1-t_n)C}{\sin C} d(x_n, y_n) + d(y_n, x_n) + d(T(x_n), y_n) \\ &\leq 2\frac{(1-t_n)C}{\sin C} s_n d(x_n, T(x_n)) + s_n d(T(x_n), x_n) \\ &+ (1-s_n) d(T(x_n), x_n) \\ &= [1 + 2\frac{(1-t_n)C}{\sin C} s_n] d(T(x_n), x_n). \end{aligned}$$

Then we have

$$t_n d(T(x_{n+1}), x_{n+1}) \leq [t_n + 2\frac{C}{\sin C} t_n(1-t_n)s_n] d(T(x_n), x_n). \tag{3.2}$$

From (3.1) and (3.2), we get

$$d(T(x_{n+1}), x_{n+1}) \leq [1 + 4\frac{C}{\sin C} t_n(1-t_n)s_n] d(T(x_n), x_n).$$

□

Lemma 15. *Let* X *be a complete* $\mathrm{CAT}(1)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$. *Suppose that* $\{t_n\}$ *and* $\{s_n\}$ *satisfy that*

$$\sum_{n=0}^{\infty} t_n(1-t_n) = \infty \quad \textit{and} \quad \sum_{n=0}^{\infty} t_n(1-t_n)s_n < \infty.$$

If $\{x_n\}$ *is defined by* (1.1) *for* $x_0 \in X$ *such that* $d(x_0, F) \leq \pi/4$, *then*

$$\lim_{n\to\infty} d(T(x_n), x_n) = 0.$$

Proof. By Lemma 11 and 14, $\lim_{n\to\infty} d(T(x_n), x_n)$ exists. Let p be the unique point in F such that $d(x_0, p) = d(x_0, F)$. By Lemma 5, we get

$$\begin{aligned} d^2(p, x_{n+1}) &= d^2(p, t_n T(y_n) \oplus (1-t_n)x_n) \\ &\leq t_n d^2(p, T(y_n)) + (1-t_n) d^2(p, x_n) - \frac{k}{2} t_n(1-t_n) d^2(T(y_n), x_n). \end{aligned}$$

Then since T is nonexpansive, we have

$$d^2(p, x_{n+1}) \leq t_n d^2(p, y_n) + (1-t_n) d^2(p, x_n) - \frac{k}{2} t_n(1-t_n) d^2(T(y_n), x_n). \tag{3.3}$$

By Lemma 5, also we get

$$\begin{aligned} d^2(p, y_n) &= d^2(p, s_n T(x_n) \oplus (1-s_n)x_n) \\ &\leq s_n d^2(p, T(x_n)) + (1-s_n) d^2(p, x_n) - \frac{k}{2} s_n(1-s_n) d^2(T(x_n), x_n). \end{aligned}$$

Since T is nonexpansive,

$$\begin{aligned} d^2(p, y_n) &\leq s_n d^2(p, x_n) + (1 - s_n) d^2(p, x_n) - \frac{k}{2} s_n (1 - s_n) d^2(T(x_n), x_n) \\ &\leq d^2(p, x_n). \end{aligned} \tag{3.4}$$

By (3.3) and (3.4), we obtain

$$d^2(p, x_{n+1}) \leq d^2(p, x_n) - \frac{k}{2} t_n (1 - t_n) d^2(T(y_n), x_n).$$

This implies

$$\frac{k}{2} \sum_{n=0}^{\infty} t_n (1 - t_n) d^2(T(y_n), x_n) \leq d^2(p, x_0) < \infty. \tag{3.5}$$

Since $\sum_{n=0}^{\infty} t_n(1 - t_n) s_n < \infty$, from (3.5), we get

$$\sum_{n=0}^{\infty} t_n (1 - t_n)[d^2(T(y_n), x_n) + s_n] < \infty.$$

Since $\sum_{n=0}^{\infty} t_n(1 - t_n) = \infty$, it implies

$$\liminf_{n\to\infty} [d^2(T(y_n), x_n) + s_n] = 0.$$

Then there exists a subsequence $\{n_k\}$ of $\{n\}$ such that

$$\lim_{k\to\infty} d(T(y_{n_k}), x_{n_k}) = 0 \quad \text{and} \quad \lim_{k\to\infty} s_{n_k} = 0. \tag{3.6}$$

Also,

$$\begin{aligned} d(T(x_{n_k}), x_{n_k}) &\leq d(T(x_{n_k}), T(y_{n_k})) + d(T(y_{n_k}), x_{n_k}) \\ &\leq d(x_{n_k}, y_{n_k}) + d(T(y_{n_k}), x_{n_k}) \\ &= s_{n_k} d(T(x_{n_k}), x_{n_k}) + d(T(y_{n_k}), x_{n_k}). \end{aligned}$$

Then it becomes

$$(1 - s_{n_k}) d(T(x_{n_k}), x_{n_k}) \leq d(T(y_{n_k}), x_{n_k}). \tag{3.7}$$

From (3.6) and (3.7), we get

$$\lim_{k\to\infty} d(T(x_{n_k}), x_{n_k}) = 0. \tag{3.8}$$

Since $\lim_{n\to\infty} d(T(x_n), x_n)$ exists, (3.8) implies that

$$\lim_{n\to\infty} d(T(x_n), x_n) = 0.$$

□

We will prove Theorem 16, which is obtained by following a first part of the proof in [5, Theorem 3.1].

Theorem 16. *Let* X *be a complete* $\mathrm{CAT}(K)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$. *Suppose that* $\{x_n\}$ *is defined by* (1.1) *under the conditions*

$$\sum_{n=0}^{\infty} t_n(1 - t_n) = \infty \ \textit{ and } \ \sum_{n=0}^{\infty} t_n(1 - t_n) s_n < \infty.$$

Then, for each $x_0 \in X$ *with* $d(x_0, F) < D_K/4$, *the sequence* $\{x_n\}$ Δ*-converges to a point of* F.

Proof. Rescaling the metric by $1/\sqrt{K}$, we may assume that $K = 1$.

Set $F_0 := F \cap B_{\pi/2}(x_0)$. For any $q \in F_0$, since $d(T(x_0), q) \leq d(x_0, q)$ and since the open ball $B_r(q)$ in X with radius $r < \pi/2$ is convex, we have

$$d(y_0, q) = d(s_0 T(x_0) \oplus (1 - s_0)x_0, q) \leq d(x_0, q).$$

Similarly, since $d(T(y_0), q) \leq d(y_0, q)$ and since the open ball $B_r(q)$ is convex, we have

$$d(x_1, q) = d(t_0 T(y_0) \oplus (1 - t_0)x_0, q) \leq d(x_0, q).$$

Using mathematical induction, we can easily get that

$$d(x_{n+1}, q) \leq d(x_n, q) \leq d(x_0, q)$$

for all $n \geq 0$. Therefore the sequence $\{x_n\}$ is Fejér monotone with respect to F_0.

Let p be the unique point in F such that $d(x_0, p) = d(x_0, F)$. Then $p \in F_0$. Also we get

$$d(x_{n+1}, p) \leq d(x_n, p) \leq d(x_0, p) < \pi/4 \tag{3.9}$$

for all $n \geq 0$. This means that the asymptotic radius $r(\{x_n\})$ of $\{x_n\}$ is less than $\pi/4$.

By Lemma 10, we only need to show that for each point x such that there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ which Δ-converges to x, it belongs to F_0. From (3.9), note that $r(p, \{x_n\}) \leq d(x_0, p) < \pi/4$.

By Lemma 7, we obtain

$$d(x, x_0) \leq d(x, p) + d(x_0, p) \leq \liminf_k d(x_{n_k}, p) + d(x_0, p) < \pi/2,$$

that is,

$$x \in B_{\pi/2}(x_0). \tag{3.10}$$

By Lemma 15, we have

$$\begin{aligned}\limsup_k d(T(x), x_{n_k}) &\leq \limsup_k d(T(x), T(x_{n_k})) + \limsup_k d(T(x_{n_k}), x_{n_k}) \\ &\leq \limsup_k d(x, x_{n_k}).\end{aligned}$$

This implies that $T(x) \in A(\{x_{n_k}\})$ and $T(x) = x$. Therefore $x \in F$. With (3.10), x belongs to F_0. By Lemma 10, it is proved. □

Since the following corollary is a direct result by letting $K = 0$ in Theorem 16, Theorem 16 is the extended result of the corresponding one in [9].

Corollary 17. *Let* X *be a complete* $\mathrm{CAT}(0)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$. *Suppose that* $\{x_n\}$ *is defined by* (1.1) *under the conditions*

$$\sum_{n=0}^{\infty} t_n(1 - t_n) = \infty \ \text{ and } \ \sum_{n=0}^{\infty} t_n(1 - t_n)s_n < \infty.$$

Then, for each $x_0 \in X$, *the sequence* $\{x_n\}$ Δ*-converges to a point of* F.

For the conditions $\sum_{n=0}^{\infty} t_n(1-t_n) = \infty$, $\sum_{n=0}^{\infty}(1-t_n)s_n < \infty$ and $\limsup_n s_n < 1$, we will get the following lemma, which is an analog of Lemma 2.12 in [2].

Lemma 18. *Let* X *be a complete* $\mathrm{CAT}(1)$ *space and let* $T : X \to X$ *be a nonexpansive mapping such that* $F := \mathrm{Fix}(T) \neq \emptyset$*. Suppose that* $\{t_n\}$ *and* $\{s_n\}$ *satisfy that*

$$\sum_{n=0}^{\infty} t_n(1-t_n) = \infty, \sum_{n=0}^{\infty} (1-t_n)s_n < \infty \text{ and } \limsup_n s_n < 1.$$

If $\{x_n\}$ *is defined by* (1.1) *for* $x_0 \in X$ *such that* $d(x_0, F) < \pi/4$*, then*

$$\lim_{n\to\infty} d(T(x_n), x_n) = 0.$$

Proof. From Equation (3.5) in Lemma 15,

$$\frac{k}{2}\sum_{n=0}^{\infty} t_n(1-t_n) d^2(T(y_n), x_n) \leq d^2(p, x_0) < \infty$$

for the unique point p in F such that $d(x_0, p) = d(x_0, F)$. Since $\sum_{n=0}^{\infty} t_n(1-t_n)$ diverges, it means that

$$\liminf_n d^2(T(y_n), x_n) = 0$$

and then

$$\liminf_n d(T(y_n), x_n) = 0. \tag{3.11}$$

Since T is nonexpansive and $d(x_n, y_n) = s_n d(T(x_n), x_n)$,

$$\begin{aligned} d(T(x_n), x_n) &\leq d(T(x_n), T(y_n)) + d(T(y_n), x_n) \\ &\leq d(x_n, y_n) + d(T(y_n), x_n) \\ &= s_n d(T(x_n), x_n) + d(T(y_n), x_n). \end{aligned}$$

Then we get

$$d(T(x_n), x_n) \leq \frac{1}{1-s_n} d(T(y_n), x_n).$$

By (3.11),

$$\liminf_n d(T(x_n), x_n) = 0. \tag{3.12}$$

Since $d(x_0, F) < \pi/4$, by Lemma 13, x_n and $T(y_n)$ are in the open ball centered at $T(x_{n+1})$ with radius $< \pi/2$. Since this open ball is convex, we have

$$\begin{aligned} d(T(x_{n+1}), x_{n+1}) &\leq t_n d(T(x_{n+1}), T(y_n)) + (1-t_n) d(T(x_{n+1}), x_n) \\ &\leq t_n d(x_{n+1}, y_n) + (1-t_n)[d(T(x_{n+1}), x_{n+1}) + d(x_{n+1}, x_n)] \\ &\leq t_n d(x_{n+1}, y_n) + (1-t_n)[d(T(x_{n+1}), x_{n+1}) + t_n d(T(y_n), x_n)]. \end{aligned}$$

Dividing by t_n, this becomes

$$d(T(x_{n+1}), x_{n+1}) \leq d(x_{n+1}, y_n) + (1-t_n) d(T(y_n), x_n).$$

By Lemma 13, $T(y_n)$ and x_n are in the open ball centered at y_n with radius $< \pi/2$, which is convex. This yields

$$d(T(x_{n+1}), x_{n+1}) \leq t_n d(T(y_n), y_n) + (1-t_n) d(x_n, y_n) + (1-t_n) d(T(y_n), x_n).$$

Since $T(x_n)$ and x_n are in the open ball centered at $T(y_n)$ with radius $< \pi/2$, we have

$$\begin{aligned} d(T(x_{n+1}), x_{n+1}) \leq{} & t_n[s_n d(T(y_n), T(x_n)) + (1-s_n) d(T(y_n), x_n)] \\ & + (1-t_n) d(x_n, y_n) + (1-t_n) d(T(y_n), x_n). \end{aligned}$$

Since T is nonexpansive and $d(x_n, y_n) = s_n d(T(x_n), x_n)$, we get

$$\begin{aligned} d(T(x_{n+1}), x_{n+1}) &\leq (1 - t_n + t_n s_n) d(x_n, y_n) + (1 - t_n s_n) d(T(y_n), x_n) \\ &\leq s_n(1 - t_n + t_n s_n) d(x_n, T(x_n)) \\ &\quad + (1 - t_n s_n)[d(T(y_n), T(x_n)) + d(T(x_n), x_n)] \\ &\leq [s_n(1 - t_n + t_n s_n) + (1 - t_n s_n)(1 + s_n)] d(x_n, T(x_n)) \\ &= [1 + 2s_n(1 - t_n)] d(x_n, T(x_n)). \end{aligned}$$

Then we get the following inequality

$$d(T(x_{n+1}), x_{n+1}) \leq [1 + 2s_n(1 - t_n)] d(x_n, T(x_n)). \tag{3.13}$$

Since $\sum s_n(1 - t_n)$ converges, applying Lemma 11 to (3.13), $\lim_{n\to\infty} d(T(x_n), x_n)$ exists. By (3.12), it is equal to zero. $\square$

By following the same proof of Theorem 16 and using Lemma 18, we obtain

Theorem 19. *Let X be a complete $\mathrm{CAT}(K)$ space and let $T : X \to X$ be a nonexpansive mapping such that $F := \mathrm{Fix}(T) \neq \emptyset$. Suppose that $\{x_n\}$ is defined by (1.1) under the conditions*

$$\sum_{n=0}^{\infty} t_n(1 - t_n) = \infty, \sum_{n=0}^{\infty} (1 - t_n)s_n < \infty \text{ and } \limsup_n s_n < 1.$$

Then, for each $x_0 \in X$ with $d(x_0, F) < D_K/4$, the sequence $\{x_n\}$ Δ-converges to a point of F.

ACKNOWLEDGEMENTS

The author would like to express his gratitude to Prof. R. Ghrist for his support. He gratefully acknowledge support from the ONR Antidote MURI project, grant no. N00014-09-1-1031.

REFERENCES

[1] M. Bridson and A. Haeflinger, Metric Spaces of Non-positive Curvature, Springer-Verlag, 1999.

[2] S. Dhompongsa and B. Panyanak, *On $\Delta-$covergence theorems in $\mathrm{CAT}(0)$ spaces*, Comput. Math. Appl. 56 (2008) 2572-2579.

[3] R. Espínola and A. Fernández-León, *$\mathrm{CAT}(K)$ spaces, weak convergence and fixed points*, J. Math. Anal. Appl. 353 (2009) 410-427.

[4] K. Goebel and W. A. Kirk, *Iteration processes for nonexpansive mappings*, in: Topological Methods in Nonlinear Functional Analysis (Toronto, Ont., 1982), in: Contemp. Math., 21, Amer. Math. Soc., Rhode Island (1983) 115-123.

[5] J. S. He, D. H. Fang, G. López and C. Li, *Mann's algorithm for nonexpansive mappings in $\mathrm{CAT}(K)$ spaces*, Nonlinear Anal. 75 (2012) 445-452.

[6] W. A. Kirk, *Geodesic geometry and fixed point theory*, in: Seminar of Mathematical Analysis (Malaga/Seville, 2002/2003), in: Colecc. Abierta, vol. 64, Univ. Sevilla Secr. Publ., Seville, (2003) 195-225.

[7] W. A. Kirk, *Geodesic geometry and fixed point theory II*, in: J. García-Falset, E. Llorens-Fuster, B. Sims (Eds.), Fixed point Theory and Applications, Yokohama Publ., (2004) 113-142.

[8] W. R. Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. 4 (1953) 506-510.

[9] B. Panyanak and T. Laokul, *On the Ishikawa iteration process in $\mathrm{CAT}(0)$ spaces*, Bull. Iranian Math. Soc. 37 (2011), no. 4, 185-197.

[10] B. Piatek, *Halpern Iteration in $\mathrm{CAT}(K)$ spaces*, Acta Math. Sinica, 27 (2011) 635-646.

[11] S. Ohta, *Convexities of metric spaces*, Geom. Dedicata, 125 (2007) 225-250.

[12] H. Zhou, *Non-expansive mappings and iterative methods in uniformly convex Banach spaces*, Acta Math. Sinica, 20 (2004) 829-836.

[13] H. Zhou, R. P. Agarwal, Y. J. Cho and Y. S. Kim, *Nonexpansive mappings and iterative methods in uniformly convex Banach spaces*, Georgian Math. J. 9 (2002) 591-600.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA PA, 19104
E-mail address: `cjun@math.upenn.edu`